

**BONAMIA EXITIOSA AND ITS INFECTION OF *CRASSOSTREA VIRGINICA*
 IN THE EASTERN USA: AN ADVISORY**

September 11, 2013

The last 14 months have seen the first two reports of the parasite *Bonamia exitiosa* infecting the eastern oyster, *Crassostrea virginica*. Hatchery-produced seed oysters from Core Sound, North Carolina were found to be infected in July 2012, and cultured seed from Stage Harbor, Chatham, Massachusetts collected in June 2013 were infected with what is presumed to be the same parasite. While *B. exitiosa* does not appear to be a serious pathogen of *C. virginica*, its emergence in *C. virginica* warrants attention from pathologists, resource managers, and the aquaculture community. Here is what we presently know about this organism.

Who is *Bonamia exitiosa*? *B. exitiosa* is a protistan parasite infecting oyster tissues. It was first observed in the oyster *Ostrea chilensis* from New Zealand, in which it is a serious pathogen. Because of the damage it causes in some hosts, *B. exitiosa* has been placed on the World Organisation for Animal Health (OIE) list of notifiable pathogens (<http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2013/>).

Distribution. While *B. exitiosa* was originally described from New Zealand¹, it is now known to occur widely, in Australia, South America, eastern and western North America, North Africa, and Europe. Along the Atlantic coast of the USA south of Cape Hatteras, the non-commercial crested oyster *Ostrea stentina* (= *Ostreola equestris*) is a host^{2,3}. The European flat oyster *Ostrea edulis*, which is established in New England, is infected by *B. exitiosa* in some European systems⁴.

Effects on oyster hosts. Parasites in the genus *Bonamia* are often referred to as “microcell haplosporidians”⁵, which emphasizes their distinctiveness from familiar haplosporidians like *Haplosporidium nelsoni* (MSX) and *H. costale* (SSO). They are tiny parasites (2-3 µm) that specifically infect and proliferate in oyster hemocytes, or blood cells. Transmission occurs directly from oyster to oyster (unlike MSX and SSO) and infections produce inflammation that, in advanced cases, can be highly disruptive of host tissue structure and function. Associated mortality in susceptible hosts like *Crassostrea ariakensis* can exceed 90%⁶. Only small *C. virginica* seed (< 20 mm) have been found infected, and while




Fig. 1. Heavy *Bonamia exitiosa* infection of *Crassostrea virginica* from North Carolina. **A.** Inflammation of oyster tissues as viewed histologically. **B.** *In situ* hybridization to the same section showing abundant *B. exitiosa* cells (dark spots) associated with oyster hemocytes. Scale bars = 50 µm. Images: N. Stokes.

prevalence of *B. exitiosa* in *C. virginica* may be high, as in the North Carolina case (93.8%), most infections have been light, and no mortality has been reported.

Environmental influences. Research on *B. exitiosa* in *C. ariakensis* indicated that the parasite was limited by salinities under 20 ppt^{7,8,9}. Based on its infection of oysters in cool temperate as well as sub-tropical systems, it must be viewed as tolerating a wide temperature range.

Detection. Due to its small size, *B. exitiosa* is difficult to detect microscopically, especially when infections are light. Use of a polymerase chain reaction assay (PCR) assay¹⁰ is recommended for general surveillance. *In situ* hybridization³ may be used to confirm infection. Control material is available from the VIMS Shellfish Pathology Laboratory. Because even the newer small-subunit ribosomal DNA-based PCR assays above cannot distinguish *B. exitiosa* from other closely related parasites, DNA sequencing of internal transcribed spacer (ITS) region sequences (still pending for the Massachusetts case) is necessary for definitive identification of *B. exitiosa* in new hosts and locations.

Management. We recommend that screening for *B. exitiosa* (preferably by PCR) be conducted in *C. virginica* seed and broodstock destined for interstate transfer, particularly when originating from the Southeast or New England, and that oysters positive for *B. exitiosa* not be transferred to areas free of the parasite. Determination of the distribution of *B. exitiosa* in wild and cultured *C. virginica* populations in the eastern USA should be a priority.

Ryan B. Carnegie, PhD
Virginia Institute of Marine Science
College of William & Mary

carnegie@vims.edu
Phone 804-684-7713

David Bushek, PhD
Haskin Shellfish Research Laboratory
Rutgers University

bushek@hsrl.rutgers.edu
Phone 856-785-0074 x4327

References

- ¹Hine PM, Cochenne-Laureau N, Berthe FC (2001) *Bonamia exitiosus* n. sp. (Haplosporidia) infecting flat oysters *Ostrea chilensis* in New Zealand. *Dis Aquat Organ* 47:63-72
- ²Carnegie RB, Barber BJ, Culloty SC, Figueras AJ, Distel DL (2000) Development of a PCR assay for detection of the oyster pathogen *Bonamia ostreae* and support for its inclusion in the Haplosporidia. *Dis Aquat Organ* 42:199-206
- ³Hill KM, Carnegie RB, Aloui-Bejaoui N, El Gharsalli R, White DM, Stokes NA, Burrenson EM (2010) Observation of a *Bonamia* sp. infecting the oyster *Ostrea stentina* in Tunisia, and a consideration of its phylogenetic affinities. *Journal of Invertebrate Pathology* 103:179-185
- ⁴Carrasco N, Villalba A, Andree KB, Engelsma MY, Lacuesta B, Ramilo A, Gairín I, Furones MD (2012) *Bonamia exitiosa* (Haplosporidia) observed infecting the European flat oyster *Ostrea edulis* cultured on the Spanish Mediterranean coast. *J Invert Path* 110:307-313
- ⁵Carnegie RB, Cochenne-Laureau N (2004) Microcell parasites of oysters: Recent insights and future trends. *Aquat Living Resour* 17:519-528
- ⁶Burrenson EM, Stokes NA, Carnegie RB, Bishop MJ (2004) *Bonamia* sp. (Haplosporidia) Found in Nonnative Oysters *Crassostrea ariakensis* in Bogue Sound, North Carolina. *J Aquat Anim Health* 16:1-9
- ⁷Bishop MJ, Carnegie RB, Stokes NA, Peterson CH, Burrenson EM (2006) Predicting the outcome of a non-native oyster introduction: facilitation of an enzootic parasite. *Mar Ecol Prog Ser* 325:145-152
- ⁸Audemard C, Carnegie RB, Bishop MJ, Peterson CH, Burrenson EM (2008) Interacting effects of temperature and salinity on *Bonamia* sp. parasitism in the Asian oyster *Crassostrea ariakensis*. *J Invert Pathol* 98:344-350
- ⁹Audemard C, Carnegie RB, Stokes NA, Bishop MJ, Peterson CH, Burrenson EM (2008) Effects of salinity on *Bonamia* sp. survival in the Asian oyster *Crassostrea ariakensis*. *Journal of Shellfish Research* 27:535-540
- ¹⁰ Carnegie RB, Stokes NA, Audemard C, Bishop MJ, Wilbur AE, Alphin TD, Posey MH, Peterson CH, Burrenson EM (2008) Strong seasonality of *Bonamia* sp. infection and induced *Crassostrea ariakensis* mortality in Bogue and Masonboro Sounds, North Carolina, USA. *J Invert Pathol* 98:335-343